

Journal of Neurotherapy: Investigations in Neuromodulation, Neurofeedback and Applied Neuroscience

News from Other Journals and Websites

Martijn Arns Senior Editor MSc
Published online: 18 May 2010.


To cite this article: Martijn Arns Senior Editor MSc (2010) News from Other Journals and Websites, Journal of Neurotherapy: Investigations in Neuromodulation, Neurofeedback and Applied Neuroscience, 14:2, 156-159, DOI: [10.1080/10874201003776248](https://doi.org/10.1080/10874201003776248)

To link to this article: <http://dx.doi.org/10.1080/10874201003776248>

PLEASE SCROLL DOWN FOR ARTICLE

© International Society for Neurofeedback and Research (ISNR), all rights reserved. This article (the “Article”) may be accessed online from ISNR at no charge. The Article may be viewed online, stored in electronic or physical form, or archived for research, teaching, and private study purposes. The Article may be archived in public libraries or university libraries at the direction of said public library or university library. Any other reproduction of the Article for redistribution, sale, resale, loan, sublicensing, systematic supply, or other distribution, including both physical and electronic reproduction for such purposes, is expressly forbidden. Preparing or reproducing derivative works of this article is expressly forbidden. ISNR makes no representation or warranty as to the accuracy or completeness of any content in the Article. From 1995 to 2013 the *Journal of Neurotherapy* was the official publication of ISNR (www.isnr.org); on April 27, 2016 ISNR acquired the journal from Taylor & Francis Group, LLC. In 2014, ISNR established its official open-access journal *NeuroRegulation* (ISSN: 2373-0587; www.neuroregulation.org).

THIS OPEN-ACCESS CONTENT MADE POSSIBLE BY THESE GENEROUS SPONSORS



NEWS FROM OTHER JOURNALS AND WEBSITES

Martijn Arns, MSc, Senior Editor

In the following section, interesting new articles recently published are summarized. The articles are mainly from the broad area of Applied Neurosciences with a focus on neurofeedback, brain computer interface, quantitative EEG, and repetitive transcranial magnetic stimulation. These are the articles, which could be found between December 1, 2009, and March 10, 2010.

Tye, K. M., Tye, L. D., Cone, J. J., Hekkelman, E. F., Janak, P. H., & Bonci, A. (2010). Methylphenidate facilitates learning-induced amygdala plasticity. *Nature Neuroscience*. Advance online publication. doi:10.1038=nn.2506

Investigating the working mechanism of methylphenidate further and showing that dopamine receptor subtypes play a role in mediating methylphenidate-induced enhancements of neural transmission and learning performance.

Hegerl, U., Himmerich, H., Engmann, B., & Hensch, T. (2010). Mania and attention-deficit=hyperactivity disorder: Common symptomatology, common pathophysiology and common treatment? *Current Opinion in Psychiatry*, 23(1), 1–7.

Review article addressing the similarities in pathophysiology in attention deficit=hyperactivity disorder and bipolar disorders, also further explaining the valuable EEG Vigilance model, which explains EEG findings for both disorders.

Schoenknecht, P., Olbrich, S., Sander, C., Spindler, P., & Hegerl, U. (2010). Treatment of acute mania with modafinil monotherapy. *Biological Psychiatry*.

Advance online publication. doi:10.1016=j.biopsych.2009.12.021

Authors employing the EEG Vigilance model to prescribe stimulant medication to a manic patient (modafenil), showing dramatic improvements in manic symptomatology as predicted from the EEG.

Nagai, Y., Cavanna, A., & Critchley, H. D. (2009). Influence of sympathetic autonomic arousal on tics: Implications for a therapeutic behavioral intervention for tourette syndrome. *Journal of Psychosomatic Research*, 67, 599–605.

GSR Biofeedback in Tourette syndrome showing that relaxation biofeedback is related to a reduction in tics, and tic frequency is positively related to sympathetic arousal during arousal session.

Philippens, I. H., & Vanwersch, R. A. (2010). Neurofeedback training on sensorimotor rhythm in marmoset monkeys. *Neuroreport*, 21, 328–332.

Dutch researchers showing that the marmoset monkeys learned to increase their SMR within five sessions, providing a further scientific basis to neurofeedback.

Helps, S. K., Broyd, S. J., James, C. J., Karl, A., Chen, W., & Sonuga-Barke, E. J. (2010). Altered spontaneous low frequency brain activity in attention deficit=hyperactivity disorder. *Brain Research*, 1322, 134–143.

Study showing that children with attention deficit=hyperactivity disorder (ADHD) showed less EEG power in very low EEG frequency bands (i.e., .02–.2 Hz). ADHD children may have deficits in maintaining a

resting brain when needed and “protecting” an active brain from intrusion of resting state brain activity. Please note these authors speak of DC-EEG, which requires a “real DC amplifier” and appropriate methodology.

Williams, L. M., Hermens, D. F., Thein, T., Clark, C. R., Cooper, N. J., Clarke, S. D., et al. (2010). Using brain-based cognitive measures to support clinical decisions in ADHD. *Pediatric Neurology*, 42(2), 118–126.

Integrative Neuroscience framework used to assess cognition, EEG and ERP measures in a large group of children with attention deficit=hyperactivity disorder ($N=175$) and a matched control group.

Stokes, D. A., & Lappin, M. S. (2010). Neurofeedback and biofeedback with 37 migraineurs: A clinical outcome study. *Behavioral and Brain Functions: BBF*, 6(1), 9.

Study demonstrating combined neuro- and biofeedback interventions were effective in reducing the frequency of migraines with clients using medication resulting in a more favorable outcome (70% experiencing at least a 50% reduction in headaches) than just medication alone (50% experience a 50% reduction) showing a large effect size ($ES=1.09$) as compared to thermal biofeedback alone ($ES=0.5$).

Paquette, V., Beaugregard, M., & Beaulieu-Pre'vost, D. (2009). Effect of a psychoneurotherapy on brain electromagnetic tomography in individuals with major depressive disorder. *Psychiatry Research*, 174, 231–239.

Study investigating psycho-neurotherapy (combination of psychotherapy and neurofeedback) in Depression, showing that clinical remission was associated with a normalization of EEG (mainly normalization of high-beta).

Ros, T., Munneke, M. A. M., Ruge, D., Gruzelier, J. H., & Rothwell, J. C. (2010). Endogenous control of waking brain rhythms induces neuroplasticity in

humans. *European Journal of Neuroscience*, 31, 770–778.

The results of this study demonstrate that a single session of alpha neurofeedback (downtraining) can produce robust increases in corticospinal excitability and decreases in intracortical inhibition of up to 150%, which last for at least 20 min.

Mulligan, B. P., Hunter, M. D., & Persinger, M. A. (2009). Effects of geomagnetic activity and atmospheric power variations on quantitative measures of brain activity: Replication of the azerbaijani studies. *Advances in Space Research*, 45, 940–948.

For the interested reader: relations between geomagnetic activity and the QEEG. Changes in right frontal theta and gamma power correlated to geomagnetic activity.

Grin-Yatsenko, V. A., Baas, I., Ponomarev, V. A., & Kropotov, J. D. (2009). Independent component approach to the analysis of EEG recordings at early stages of depressive disorders. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, 121, 281–289.

Study investigating Independent Components Analysis and sLORETA methods in early stages of depression.

Hunter, A. M., Muthe'n, B. O., Cook, I. A., & Leuchter, A. F. (2010). Antidepressant response trajectories and quantitative electroencephalography (QEEG) biomarkers in major depressive disorder. *Journal of Psychiatric Research*, 44, 90–98.

Investigating the effects of a volatile course of symptom change in depression, which exert a confounding effect on typical binary endpoint outcomes. The relation to EEG concordance is discussed.

Velikova, S., Locatelli, M., Insacco, C., Smeraldi, E., Comi, G., & Leocani, L. (2010). Dysfunctional brain circuitry in obsessive-compulsive disorder: Source and coherence analysis of EEG rhythms. *Neuroimage*, 49(1), 977–983.

In obsessive-compulsive disorder, increased frontal beta is consistent with previous evidence of frontal dysfunction. Hyperactivity of insular delta sources, together with rhythms decoupling and reduced interhemispheric alpha coherence, are consistent with additional involvement of cortico-subcortical functional connections. Combined use of power and coherence analysis may provide functional measures on different levels of involvement of cortico-subcortical circuits in neuropsychiatric disorders.

Fedotchev, A. I. (2010). Efficacy of EEG biofeedback procedures in correcting stress-related functional disorders. *Human Physiology*, 36, 86–90.

Description of neurofeedback approaches to stress-related disorders employing “double EEG Biofeedback” or two independent feedback loops: the traditional feedback loop and an additional resonance stimulation loop.

Narushima, K., McCormick, L. M., Yamada, T., Thatcher, R. W., & Robinson, R. G. (2010). Subgenual cingulate theta activity predicts treatment response of repetitive transcranial magnetic stimulation in participants with vascular depression. *Journal of Neuropsychiatry and Clinical Neurosciences*, 22, 75.

Increased low-theta power in the subgenual anterior cingulate cortex predicts response to repetitive transcranial magnetic stimulation.

Rothkegel, H., Sommer, M., & Paulus, W. (2010). Breaks during 5 hz rTMS are essential for facilitatory after effects. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, 121, 426–430.

This study demonstrates that breaks during high-frequency transcranial magnetic stimulation protocols are required for plastic changes to occur as opposed to continuous stimulation.

Chistyakov, A. V., Rubicsek, O., Kaplan, B., Zaaroor, M., & Klein, E. (2010). Safety, tolerability and preliminary evidence for

antidepressant efficacy of theta-burst transcranial magnetic stimulation in patients with major depression. *The International Journal of Neuropsychopharmacology= Official Scientific Journal of the Collegium Internationale Neuropsychopharmacologicum (CINP)*, 1–7.

Study demonstrating that Theta burst stimulation (repetitive transcranial magnetic stimulation) is safe and well tolerated in depressed patients and seems to have antidepressant properties.

Schutter, D. J. (2010). Quantitative review of the efficacy of slow-frequency magnetic brain stimulation in major depressive disorder. *Psychological Medicine*. Advance online publication. doi: 10.1017/S003329171000005X

Meta-analysis on 1 Hz right frontal repetitive transcranial magnetic stimulation in depression demonstrating a similar efficacy to 10 Hz Left frontal stimulation.

Minc, D., Machado, S., Bastos, V. H., Machado, D., Cunha, M., Cagy, M., et al. (2010). Gamma band oscillations under influence of bromazepam during a sensorimotor integration task: An EEG coherence study. *Neuroscience Letters*, 469, 145–149.

They propose that the gamma band plays an important role in the binding among several brain areas in complex motor tasks and that each hemisphere is influenced in a different manner by the neuromodulator.

Papageorgiou, T., Curtis, W. A., McHenry, M., & Laconte, S. M. (2009). Neurofeedback of two motor functions using supervised learning-based real-time functional magnetic resonance imaging. *Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society. Conference*, 1, 5377–5380.

This study examines the effects of neurofeedback provided by support vector machine classification-based real-time functional magnetic resonance imaging during two types of motor tasks.

Gil, Y., Li, G., & Lee, J. (2009). Integrated real-time neurofeedback system to raise the frontal lobe activity: Design and implementation. *Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society. Conference, 1*, 84-848.

In this article, an integrated portable and real-time neurofeedback system assisted by EEG has been developed. The algorithm for this system has been developed and its performance has been verified by the functional magnetic resonance imaging experiment.

Walker, J. E. (2009). Using qEEG-guided neurofeedback for epilepsy versus standardized protocols: Enhanced effectiveness? *Applied Psychophysiology and Biofeedback, 35*, 29-30.

A review of EEG treatments for epilepsy and a discussion on how QEEG might potentially enhance effectiveness.

Birbaumer, N., Ramos Murguialday, A., Weber, C., & Montoya, P. (2009). Neurofeedback and brain-computer interface clinical applications. *International Review of Neurobiology, 86*, 107-117.

A review of developments in Brain Computer Interfaces ranging from EEG based, to functional magnetic resonance imaging-based implementations.

Ehlers, C. L., Phillips, E., Gizer, I. R., Gilder, D. A., & Wilhelmsen, K. C. (2010). EEG spectral phenotypes: Heritability and association with marijuana and alcohol dependence in an American Indian community study. *Drug and Alcohol Dependence, 106*, 101-110.

Data suggest that the traits of EEG delta and high beta=gamma activity are correlated with marijuana dependence and alcohol dependence, respectively, in this community sample of Native Americans.

NEWS FROM OTHER JOURNALS AND WEBSITES

Martijn Arns, MSc, Senior Editor

In the following section, interesting new articles recently published are summarized. The articles are mainly from the broad area of Applied Neurosciences with a focus on neurofeedback, brain computer interface, quantitative EEG, and repetitive transcranial magnetic stimulation. These are the articles, which could be found between December 1, 2009, and March 10, 2010.

Tye, K. M., Tye, L. D., Cone, J. J., Hekkelman, E. F., Janak, P. H., & Bonci, A. (2010). Methylphenidate facilitates learning-induced amygdala plasticity. *Nature Neuroscience*. Advance online publication. doi:10.1038/nn.2506

Investigating the working mechanism of methylphenidate further and showing that dopamine receptor subtypes play a role in mediating methylphenidate-induced enhancements of neural transmission and learning performance.

Hegerl, U., Himmerich, H., Engmann, B., & Hensch, T. (2010). Mania and attention-deficit/hyperactivity disorder: Common symptomatology, common pathophysiology and common treatment? *Current Opinion in Psychiatry*, 23(1), 1–7.

Review article addressing the similarities in pathophysiology in attention deficit/hyperactivity disorder and bipolar disorders, also further explaining the valuable EEG Vigilance model, which explains EEG findings for both disorders.

Schoenknecht, P., Olbrich, S., Sander, C., Spindler, P., & Hegerl, U. (2010). Treatment of acute mania with modafinil monotherapy. *Biological Psychiatry*.

Advance online publication. doi:10.1016/j.biopsych.2009.12.021

Authors employing the EEG Vigilance model to prescribe stimulant medication to a manic patient (modafenil), showing dramatic improvements in manic symptomatology as predicted from the EEG.

Nagai, Y., Cavanna, A., & Critchley, H. D. (2009). Influence of sympathetic autonomic arousal on tics: Implications for a therapeutic behavioral intervention for tourette syndrome. *Journal of Psychosomatic Research*, 67, 599–605.

GSR Biofeedback in Tourette syndrome showing that relaxation biofeedback is related to a reduction in tics, and tic frequency is positively related to sympathetic arousal during arousal session.

Philippens, I. H., & Vanwersch, R. A. (2010). Neurofeedback training on sensorimotor rhythm in marmoset monkeys. *Neuroreport*, 21, 328–332.

Dutch researchers showing that the marmoset monkeys learned to increase their SMR within five sessions, providing a further scientific basis to neurofeedback.

Helps, S. K., Broyd, S. J., James, C. J., Karl, A., Chen, W., & Sonuga-Barke, E. J. (2010). Altered spontaneous low frequency brain activity in attention deficit/hyperactivity disorder. *Brain Research*, 1322, 134–143.

Study showing that children with attention deficit/hyperactivity disorder (ADHD) showed less EEG power in very low EEG frequency bands (i.e., .02–.2 Hz). ADHD children may have deficits in maintaining a

resting brain when needed and “protecting” an active brain from intrusion of resting state brain activity. Please note these authors speak of DC-EEG, which requires a “real DC amplifier” and appropriate methodology.

Williams, L. M., Hermens, D. F., Thein, T., Clark, C. R., Cooper, N. J., Clarke, S. D., et al. (2010). Using brain-based cognitive measures to support clinical decisions in ADHD. *Pediatric Neurology*, 42(2), 118–126.

Integrative Neuroscience framework used to assess cognition, EEG and ERP measures in a large group of children with attention deficit/hyperactivity disorder ($N = 175$) and a matched control group.

Stokes, D. A., & Lappin, M. S. (2010). Neurofeedback and biofeedback with 37 migraineurs: A clinical outcome study. *Behavioral and Brain Functions: BBF*, 6(1), 9.

Study demonstrating combined neuro- and biofeedback interventions were effective in reducing the frequency of migraines with clients using medication resulting in a more favorable outcome (70% experiencing at least a 50% reduction in headaches) than just medication alone (50% experience a 50% reduction) showing a large effect size ($ES = 1.09$) as compared to thermal biofeedback alone ($ES = 0.5$).

Paquette, V., Beaugard, M., & Beaulieu-Prévost, D. (2009). Effect of a psychoneurotherapy on brain electromagnetic tomography in individuals with major depressive disorder. *Psychiatry Research*, 174, 231–239.

Study investigating psycho-neurotherapy (combination of psychotherapy and neurofeedback) in Depression, showing that clinical remission was associated with a normalization of EEG (mainly normalization of high-beta).

Ros, T., Munneke, M. A. M., Ruge, D., Gruzelier, J. H., & Rothwell, J. C. (2010). Endogenous control of waking brain rhythms induces neuroplasticity in

humans. *European Journal of Neuroscience*, 31, 770–778.

The results of this study demonstrate that a single session of alpha neurofeedback (downtraining) can produce robust increases in corticospinal excitability and decreases in intracortical inhibition of up to 150%, which last for at least 20 min.

Mulligan, B. P., Hunter, M. D., & Persinger, M. A. (2009). Effects of geomagnetic activity and atmospheric power variations on quantitative measures of brain activity: Replication of the azerbaijani studies. *Advances in Space Research*, 45, 940–948.

For the interested reader: relations between geomagnetic activity and the QEEG. Changes in right frontal theta and gamma power correlated to geomagnetic activity.

Grin-Yatsenko, V. A., Baas, I., Ponomarev, V. A., & Kropotov, J. D. (2009). Independent component approach to the analysis of EEG recordings at early stages of depressive disorders. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, 121, 281–289.

Study investigating Independent Components Analysis and sLORETA methods in early stages of depression.

Hunter, A. M., Muthén, B. O., Cook, I. A., & Leuchter, A. F. (2010). Antidepressant response trajectories and quantitative electroencephalography (QEEG) biomarkers in major depressive disorder. *Journal of Psychiatric Research*, 44, 90–98.

Investigating the effects of a volatile course of symptom change in depression, which exert a confounding effect on typical binary endpoint outcomes. The relation to EEG cordance is discussed.

Velikova, S., Locatelli, M., Insacco, C., Smeraldi, E., Comi, G., & Leocani, L. (2010). Dysfunctional brain circuitry in obsessive-compulsive disorder: Source and coherence analysis of EEG rhythms. *Neuroimage*, 49(1), 977–983.

In obsessive-compulsive disorder, increased frontal beta is consistent with previous evidence of frontal dysfunction. Hyperactivity of insular delta sources, together with rhythms decoupling and reduced interhemispheric alpha coherence, are consistent with additional involvement of cortico-subcortical functional connections. Combined use of power and coherence analysis may provide functional measures on different levels of involvement of cortico-subcortical circuits in neuropsychiatric disorders.

Fedotchev, A. I. (2010). Efficacy of EEG biofeedback procedures in correcting stress-related functional disorders. *Human Physiology*, 36, 86–90.

Description of neurofeedback approaches to stress-related disorders employing “double EEG Biofeedback” or two independent feedback loops: the traditional feedback loop and an additional resonance stimulation loop.

Narushima, K., McCormick, L. M., Yamada, T., Thatcher, R. W., & Robinson, R. G. (2010). Subgenual cingulate theta activity predicts treatment response of repetitive transcranial magnetic stimulation in participants with vascular depression. *Journal of Neuropsychiatry and Clinical Neurosciences*, 22, 75.

Increased low-theta power in the subgenual anterior cingulate cortex predicts response to repetitive transcranial magnetic stimulation.

Rothkegel, H., Sommer, M., & Paulus, W. (2010). Breaks during 5 Hz rTMS are essential for facilitatory after effects. *Clinical Neurophysiology: Official Journal of the International Federation of Clinical Neurophysiology*, 121, 426–430.

This study demonstrates that breaks during high-frequency transcranial magnetic stimulation protocols are required for plastic changes to occur as opposed to continuous stimulation.

Chistyakov, A. V., Rubicsek, O., Kaplan, B., Zaaroor, M., & Klein, E. (2010). Safety, tolerability and preliminary evidence for

antidepressant efficacy of theta-burst transcranial magnetic stimulation in patients with major depression. *The International Journal of Neuropsychopharmacology/Official Scientific Journal of the Collegium Internationale Neuropsychopharmacologicum (CINP)*, 1–7.

Study demonstrating that Theta burst stimulation (repetitive transcranial magnetic stimulation) is safe and well tolerated in depressed patients and seems to have antidepressant properties.

Schutter, D. J. (2010). Quantitative review of the efficacy of slow-frequency magnetic brain stimulation in major depressive disorder. *Psychological Medicine*. Advance online publication. doi: 10.1017/S003329171000005X

Meta-analysis on 1 Hz right frontal repetitive transcranial magnetic stimulation in depression demonstrating a similar efficacy to 10 Hz Left frontal stimulation.

Minc, D., Machado, S., Bastos, V. H., Machado, D., Cunha, M., Cagy, M., et al. (2010). Gamma band oscillations under influence of bromazepam during a sensorimotor integration task: An EEG coherence study. *Neuroscience Letters*, 469, 145–149.

They propose that the gamma band plays an important role in the binding among several brain areas in complex motor tasks and that each hemisphere is influenced in a different manner by the neuromodulator.

Papageorgiou, T., Curtis, W. A., McHenry, M., & Laconte, S. M. (2009). Neurofeedback of two motor functions using supervised learning-based real-time functional magnetic resonance imaging. *Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society. Conference*, 1, 5377–5380.

This study examines the effects of neurofeedback provided by support vector machine classification-based real-time functional magnetic resonance imaging during two types of motor tasks.

Gil, Y., Li, G., & Lee, J. (2009). Integrated real-time neurofeedback system to raise the frontal lobe activity: Design and implementation. *Conference Proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society. IEEE Engineering in Medicine and Biology Society. Conference, 1*, 84–848.

In this article, an integrated portable and real-time neurofeedback system assisted by EEG has been developed. The algorithm for this system has been developed and its performance has been verified by the functional magnetic resonance imaging experiment.

Walker, J. E. (2009). Using qEEG-guided neurofeedback for epilepsy versus standardized protocols: Enhanced effectiveness? *Applied Psychophysiology and Biofeedback, 35*, 29–30.

A review of EEG treatments for epilepsy and a discussion on how QEEG might potentially enhance effectiveness.

Birbaumer, N., Ramos Murguialday, A., Weber, C., & Montoya, P. (2009). Neurofeedback and brain-computer interface clinical applications. *International Review of Neurobiology, 86*, 107–117.

A review of developments in Brain Computer Interfaces ranging from EEG based, to functional magnetic resonance imaging-based implementations.

Ehlers, C. L., Phillips, E., Gizer, I. R., Gilder, D. A., & Wilhelmsen, K. C. (2010). EEG spectral phenotypes: Heritability and association with marijuana and alcohol dependence in an American Indian community study. *Drug and Alcohol Dependence, 106*, 101–110.

Data suggest that the traits of EEG delta and high beta/gamma activity are correlated with marijuana dependence and alcohol dependence, respectively, in this community sample of Native Americans.